

Discovery of a Narrow-Line Seyfert 1 galaxy

K. P. Singh ^a

^a*Department of Astronomy & Astrophysics, Tata Institute of Fundamental Research, Mumbai, India*

L. R. Jones ^b

^b*Department of Physics & Astronomy, University of Birmingham, U.K.*

Abstract

We have discovered a Narrow-Line Seyfert 1 (NLS1) galaxy and identified it with an ultrasoft X-ray source (**RX 215319-1514**) detected with *ROSAT*. We present its X-ray and optical spectral characteristics. Its redshift is found to be 0.0778 ± 0.0002 . Its optical spectrum shows fairly strong Fe II emission lines, and its X-ray spectrum has an extremely steep power-law index of $\Gamma \geq 4$, making it an extremely interesting example of this class of AGNs. No low-energy absorption or variability is observed. The soft X-ray (0.1–2.0 keV) luminosity is estimated to be $3.4\text{--}12.5 \times 10^{43}$ ergs s^{−1}, depending on the spectral model.

Key words: galaxies: active; quasars: general; X-rays: galaxies

1 Introduction

Narrow-line Seyfert 1 (NLS1) galaxies with narrow (FWHM= 500–2000 km s^{−1}) emission lines of hydrogen in their optical/UV spectrum tend to have the steepest soft X-ray spectra in the *ROSAT* band of 0.1–2.0 keV, and form a distinct class of active galactic nuclei (see Boller et al. 1996 and references therein). Catalogues of ultra-soft X-ray sources are, therefore, very useful to find such objects. While carrying out optical spectroscopy of candidate objects in the positional error circles of such sources from the catalogue given by Singh et al. (1995), we have found a new narrow-line Seyfert 1 (NLS1) galaxy. The galaxy is a strong candidate for being a counterpart to the ultra-soft X-ray source – WGA J2153.3-1514 (RXJ 2153.3-1514). A brief summary of its optical and X-ray properties is presented below.

2 X-ray and Optical Observations

Analysis of archival *ROSAT* PSPC data in the vicinity of WGA J2153.3-1514 shows two bright optical objects near the X-ray peak – a 14.7 mag galaxy and a 11.72 mag star (see Figure 1). The X-ray beam is broadened, and somewhat distorted, due to the large off-axis angle (41 arcmin) of the source.




Fig. 1. Overlay of X-ray contour map from *ROSAT* PSPC observations on optical image from digitised sky survey.

Optical spectra were taken for both the objects at the Lick 3m telescope [Slit width = 2.5 arcsec; Grism for blue and a grating for red spectrum both of 600 lines/mm gave a resolution of $6.5 \text{ \AA}$ (FWHM)]. The spectrum of the galaxy is shown in Figure 2 after smoothing. The spectral parameters for the emission lines are given in Table 1. Spectral features are consistent with the classification as a NLS1 galaxy (Osterbrock & Pogge 1985; Goodrich 1989). The star shows a spectrum of a late-A to an early-F type, and no chromospheric emission or emission lines characteristic of an accreting cataclysmic variable are detected. These properties and the ratio of X-ray flux (F_x) to optical flux (F_v), effectively rule out the star as being the candidate for the X-ray source.

Table 1. Optical Emission Line Parameters of the NLS1 galaxy

Emission Line	Equivalent Width ($\text{\AA}$)	Measured FWHM ($\text{\AA}$)	Intrinsic FWHM (km s^{-1})
NII 6549 $\text{\AA}$	1.0 ± 0.3	1.1 ± 0.5	-
H α	40.8 ± 1.0	22.3 ± 1.2	880 ± 50
NII 6583 $\text{\AA}$	9.8 ± 0.6	7.8 ± 0.6	180 ± 25




Fig. 2. Optical spectrum (smoothed to show Fe II features) of the active galaxy.

X-ray observations from 2 epochs – 1992 and 1993, analysed separately as well as jointly for their spectral content, show a very steep spectrum ($\Gamma \geq 4$ for power-law model). Spectral parameters for the power-law and other models are given in Table 2. No variability in the X-ray flux or spectral parameters is detected. Presence of any low-energy absorption local to the source is also not detected. Galactic N_{H} in this direction is $4 \times 10^{20} \text{ cm}^{-2}$, based on 21 cm radio observations.

Table 2. X-ray Spectral Parameters

Model	$\Gamma/\text{kT}(\text{eV})$	N_{H}	F_x	S_x	L_x
		10^{20}	10^{-14}ergs	10^{-12}ergs	$10^{43}\text{ergs s}^{-1}$
		cm^{-2}	$\text{cm}^{-2}\text{s}^{-1}$	$\text{cm}^{-2}\text{s}^{-1}$	$(z=0.0778)$
					$(H_o=75; q_o=0)$
Power-law	$6.4^{+5}_{-1.5}$	$4^{+3.9}_{-1.4}$	9.5	10	12.5
Black-body	30^{+9}_{-9}	4	7.5	2.7	3.4
MEKAL	54^{+11}_{-12}	4	14	5.3	6.6

Steep X-ray spectra, as are observed here, are usually associated with NLS1 galaxies (Brandt, Pounds, & Fink 1995; Boller, Brandt, & Fink 1996; Laor et al. 1997). The position of this galaxy on the Γ - $H\beta$ line width plane (Fig. 8 of Boller et al (1996)) is consistent with an extrapolation based on the other NLS1 galaxies. For further details, please refer to Singh & Jones (2000).

Higher resolution X-ray observations with more sensitive satellites like *Chandra* and *XMM – Newton*, as well as higher resolution optical/UV spectra with better signal-to-noise ratio are required to fully understand the nature of the AGN in this galaxy.

References

- [1] T. Boller, W.N. Brandt, & H.H. Fink, *A&A* **305** (1996) 53.
- [2] W.N. Brandt, K.A. Pounds, & H.H. Fink, *MNRAS* **273** (1995) L47.
- [3] R.W. Goodrich, *ApJ* **342** (1989) 224.
- [4] A. Laor, F. Fiore, M. Elvis, B.J. Wilkes, & J.C. McDowell, *ApJ* **477** (1997) 93.
- [5] D.E. Osterbrock, & R.W. Pogge, *ApJ* **297** (1985) 166.
- [6] K.P. Singh, P. Barrett, N.E. White, P. Giommi, & L. Angelini, *ApJ* **455** (1995) 456.
- [7] K.P. Singh, & L.R. Jones, *MNRAS* **312** (2000) 380.
- [8] N.E. White, P. Giommi, & L. Angelini, IAU Circ. No. 610 (1994)